

STOP THESE VITAMINS AND MEDICATIONS AT LEAST 10 DAYS BEFORE SURGERY

Do not take any Vitamin E for 10 days before and after surgery.

Do not take any medications, which contain aspirin (whether prescription or over the counter) or other blood thinners for 10 days before surgery. Please check the labels of any medications you take to make sure they do not contain aspirin. If you are not sure, ask your doctor or pharmacist. The following is a partial list of products to avoid before surgery: **If you take coumadin or warfarin your primary care physician must decide how many days you need to be off these medications.**

Acetyl salicylic acid	Damason-P	Nabumetone
Advil	Darvon w/A.S.A.	Nalfon
Aleve	Darvon-N w/A.S.A.	Naprosyn
Alka-Seltzer Effervescent Pain Reliever	Dia-Gesic	Naproxen
Anacin (regular and maximum strength)	Diclofenac	Norgesic, Norgesic Forte
Anaprox	Disalcid	Norwich aspirin
Anodynos DIIC	Dolobid	Nuprin
Ansaid	Dolene compound	Pentasa
Apa-San	Dolprin	Percodan, Percodan-Demi
Arthritis Pain Formula	Doxaphene compound	Phentermine
Asacol	Duradyne	Plavix
A.S.A.	Easprin	Ponstel
A.S.A. Enseals	Ecotrin	Presalin
Ascriptin, Ascriptin A/D,	Emcodeine	Relafen
Ascriptin w/codeine	Empirin, Empirin w/codeine	Rid-A-Pain w/codeine
Aspergum	Equagesic	Salatin
Axotal	Equazine-M	Saleto
B-A-C, B-A-C w/codeine	Excedrin	Salfex
Bayer aspirin	Feldene	Salocol
Maximum Bayer aspirin	Fiorgen PF	Soma compound
Bayer Children's aspirin	Fiorinal, Fiorinal w/codeine	St. Joseph children's aspirin
Bayer Timed Release	Gemnisyn	Supac
Bexophene	Ibuprofen	Synalgos DC
Buffaprin	Indocin	Talwin compound
Bufferin	Isollyl Improved	Ticlid
Buffets II	Lanorinal	Tolectin
Buffex	Lodine	Toradol
Buffinol	Magnaprin	Trilisate
Butalbital compound	Maprin	Trigesic
Cama Arthritis Reliever	Marnal	Uracel
Carisoprodol	Measurin	Vanquish
Codoxy	Meclomen	Voltaren
Cope	Meprogesic Q	Warfarin
Coumadin(usually stopped 3 days prior to surgery and restarted 2 days after surgery)	Micrainin	Wesprin buffered
	Midol	Zorprin
	Motrin	
	Mono-Gesic	